

Health Expectancy in Spain

What is health expectancy?

Health expectancies were first developed to address whether or not longer life is being accompanied by an increase in the time lived in good health (the compression of morbidity scenario) or in bad health (expansion of morbidity). So health expectancies divide life expectancy into life spent in different states of health, from say good to bad health. In this way they add a dimension of quality to the quantity of life lived.

How is the effect of longer life measured?

The general model of health transitions (WHO, 1984) shows the differences between life spent in different states: total survival, disability-free survival and survival without chronic disease. This leads naturally to life expectancy (the area under the 'mortality' curve), disability-free life expectancy (the area under the 'disability' curve) and life expectancy without chronic disease (the area under the 'morbidity' curve).

The general model of health transition (WHO, 1984): observed mortality and hypothetical morbidity and disability survival curves for females, USA, 1980



e_0^{**} and e_{60}^{**} are the number of years of autonomous life expected at birth and at age 60, respectively.
 M_{50}^{**} is the age to which 50% of females could expect to survive without loss of autonomy.

There are in fact as many health expectancies as concepts of health. The commonest health expectancies are those based on self-perceived health, activities of daily living and on chronic morbidity.

How do we compare health expectancies?

Health expectancies are independent of the size of populations and of their age structure and so they allow direct comparison of different population sub-groups: e.g. sexes, socio-professional categories, as well as countries within Europe (Robine et al., 2003).

Health expectancies are most often calculated by the Sullivan method (Sullivan, 1971). However to make

valid comparisons, the underlying health measure should be truly comparable.

To address this, the European Union has decided to include a small set of health expectancies among its European Core Health Indicators (ECHI) to provide summary measures of disability (i.e., activity limitation), chronic morbidity and perceived health. Therefore the Minimum European Health Module (MEHM), composed of 3 general questions covering these dimensions, has been introduced into the Statistics on Income and Living Conditions (SILC) to improve the comparability of health expectancies between countries.* In addition life expectancy without long term activity limitation, based on the disability question, was selected in 2004 to be one of the structural indicators for assessing the EU strategic goals (Lisbon strategy) under the name of “Healthy Life Years” (HLY).

Further details on the MEHM, the European surveys and health expectancy calculation and interpretation can be found on www.eurohex.eu.

What is in this report?

This report is produced by the European Health and Life Expectancy Information System (EHLEIS) as part of a country series. In each report we present:

- Life expectancies and Healthy Life Years (HLY) at age 65 for the country of interest and for the overall 28 European Union member states (EU28), using the SILC question on long term health related disability, known as the GALI (Global Activity Limitation Indicator), from 2004 to 2014. The wording of the question has been revised in 2008;
- Prevalence of activity limitation in the country of interest and in the European Union based on the GALI question by sex and age group;
- Health expectancies based on the two additional dimensions of health (chronic morbidity and self-perceived health) for the country of interest, based on SILC 2014;
- Prevalence of activity limitation in Europe (EU28) in 2005, 2008, 2011 and 2014.

References

Jagger C., Gillies C., Moscone F., Cambois E., Van Oyen H., Nusselder W., Robine J.-M., EHLEIS Team. Inequalities in healthy life years in the 25 countries of the European Union in 2005: a cross-national meta-regression analysis. *The Lancet*. 2008;372(9656) 2124-2131
Robine J.-M., Jagger C., Mathers C.D., Crimmins E.M., Suzman R.M., Eds. *Determining health expectancies*. Chichester UK: Wiley, 2003.
Sullivan D.F. *A single index of mortality and morbidity*. HSMHA Health Reports 1971;86:347-354.
World Health Organization. *The uses of epidemiology in the study of the elderly: Report of a WHO Scientific Group on the Epidemiology of Aging*. Geneva: WHO, 1984 (Technical Report Series 706).

* Before the revision of 2008, the translations of the module used in some countries were not optimum (See Eurostat-EU Task Force on Health Expectancies common statement about the SILC data quality). The revision is being evaluated.

Life expectancy (LE) and Healthy Life Years (HLY) at age 65 for Spain and the European Union (EU28) based on SILC (2004-2014)

Key points:


Spanish life expectancy (LE) at age 65 has increased by 2.0 years for women and men over the period 2004-2014. LE was above the EU28 average (21.6 for women and 18.1 for men) in 2014.

The new HLY series, initiated in 2004 with the SILC data, shows values for Spain being in 2014 above the EU28 average (8.6 for women and men) by 0.9 year for women and 1.5 year for men.


In 2014 women and men at age 65 can expect to spend 40% and 52% of their life without *self-reported long-term activity limitations* respectively.

Note that the wording of the GALI question was changed in Spain in 2008 to better reflect the EU standard.

This may explain the strong decrease in HLY observed for men and women between 2007 and 2008. Between 2010 and 2011 HLY increased for both sexes, decreased slightly in 2012 and by 2013 rose to its previous level. Between 2013 and 2014 HLY increased notably for both sexes.




Prevalence of activity limitation in Spain and in the European Union (EU28) based on the GALI question, by sex and age group (SILC, Mean 2012-2014)



Reports of limitation in usual activities strongly increase with age in the European Union and women systematically report slightly more activity limitation than men. Compared to the mean trajectory by age observed in the European Union in the 3 years 2012-2014, Spain tends to display lower prevalence rate of activity limitation at all ages except at age 85 and over where the prevalence become slightly higher than EU28 for women only.

These results should be interpreted with caution as samples sizes in the SILC survey vary remarkably; for instance in 2014 they ranged from 5758 in Denmark to 40274 in Italy. In 2014, the sample size for Spain comprised 13840 women and 12673 men aged 16 years and over.

Life and health expectancies at age 65 based on activity limitation (Healthy Life Years), chronic morbidity and perceived health for Spain (Health data from SILC 2014)



Key points:

In 2014, LE at age 65 in Spain was 23.5 years for women and 19.3 years for men.

Based on the SILC 2014, at age 65, women spent 9.5 years (40% of their remaining life) without activity limitation (corresponding to Healthy Life Years (HLY)), 9.5 years (40%) with moderate activity limitation and 4.5 years (19%) with severe activity limitation.*

Men of the same age spent 10.1 years (52% of their remaining life) without activity limitation compared to 6.8 years (35%) with moderate activity limitation and 2.4 years (12%) with severe activity limitation.*

Although total years lived by men were less than those for women, for life expectancy in very good or good perceived health and for life expectancy without activity limitation the years of life spent in positive health were greater for men than women. Therefore compared to men, women spent a larger number of years and a larger proportion in ill health and these years of ill health were more likely to be years with severe health problems.

These results should be interpreted cautiously given the lack of the institutional population, such as people living in nursing homes.

* These may not sum to Life Expectancy due to rounding


Publications and reports on health expectancies for Spain

- Puga, M., Pujol, R. and Abellán, A. (2014). EVOLUCIÓN Y DIFERENCIAS TERRITORIALES DE LA ESPERANZA DE VIDA LIBRE DE DISCAPACIDAD A LOS 65 AÑOS EN ESPAÑA. XIV Congreso Nacional de Población de la Asociación de Geógrafos Españoles, Sevilla. https://www.researchgate.net/publication/265787613_Evolucion_y_diferencias_territoriales_de_la_Esperanza_de_Vida_Libre_de_Discapacidad_a_los_65_años_en_España
- Esperanzas de vida en España, 2013. Madrid: Ministerio de Sanidad, Servicios Sociales e Igualdad, 2015. http://www.msssi.gob.es/estadEstudios/estadisticas/inforRecopilaciones/EsperanzasDeVida_2013.pdf
- Martin U, Esnaola S. Changes in social inequalities in disability-free life expectancy in Southern Europe: the case of the Basque Country. *Int J Equity Health*. 2014 Sep 20; 13(1):74.
- Ministerio de Sanidad, Servicios Sociales e Igualdad. Indicadores de Salud 2013. Evolución de los indicadores del estado de salud en España y su magnitud en el contexto de la Unión Europea. Madrid: Ministerio de Sanidad, Servicios Sociales e Igualdad, 2014.
- Solé-Auró A, Alcañiz M. Are we living longer but less healthy? Trends in mortality and morbidity in Catalonia (Spain), 1994-2011. *Eur J Ageing* 2014 DOI 10.1007/s10433-014-0317-9.
- Gutiérrez-Fisac JL, Suárez M, Neira M, Regidor E. Esperanzas de vida en salud en España 2007-2011. Años de vida saludable en España y sus comunidades autónomas. Madrid: Ministerio de Sanidad, Servicios Sociales e Igualdad, 2013.
- Gutiérrez-Fisac J.L., Regidor E., Alfaro M. *Healthy Life Expectancies in Spain 1986-2007. Disability-free life expectancy and Life expectancy in good perceived health in Spain and its Autonomous Communities*. Madrid: Ministry of Health, Social Policy and Equality, 2010.
- Morata L.B., Losilla C.B., Estany M.G., Gondolbeu K.M. Tipologías sociodemográficas de individuos con dependencia en España y supervivencia en estado de Salud [Sociodemographic typologies of individuals with dependence in Spain and healthy life expectancy]. *Rev Esp Geriatr Gerontol*. 2008; 43(1):19-31.
- Jagger C., Gillies C., Mascone F., Cambois E., Van Oyen H., Nusselder W.J., Robine J.-M., EHLEIS team. Inequalities in healthy life years in the 25 countries of the European Union in 2005: a cross-national meta-regression analysis. *The Lancet*. 2008; 372(9656):2124-2131.
- Gispert R., Ruiz-Ramos M., Bares M.A., Viciano F., Clot-Razquin G. Differences in disability-free life expectancy by gender and autonomous regions in Spain [Differences in Disability-Free Life Expectancy by Gender and Autonomous Regions in Spain]. *Rev Esp Salud Pública*. 2007 Mar-Apr; 81(2):155-165.

Prevalence of activity limitation in Europe (EU28) in 2005, 2008, 2011 and 2014

Thanks to the EU-SILC survey, we now have 10 years of experience in measuring disability within the European Union. The survey really started in 2005 with 25 Member States (MS). In 2008, a coordinated revision of the translation of the GALI was made by some countries to better reflect the original standard. An evaluation made by Eurostat shows that in 2012 the translation of the GALI fully follows the English standard in 18 MS, partially in 8 others and still not in 5 MS. Progressively EU-SILC involved 27 then 28 MS but all the estimations provided below have been estimated for the EU28. The prevalence of disability among women and men is displayed by age and level of severity of the reported disability, from the age group 16-19 to 85+, for the calendar years 2005, 2008, 2011 and 2014.

Prevalence of activity limitation in Europe (EU28), by sex and age group, SILC EU28, 2005, 2008, 2011 and 2014



The revision of the translation of the GALI in 2008 significantly changed the age trajectory of the prevalence of disability, increasing the report of disability among the oldest participants in the EU-SILC survey, especially for those reporting being not severely limited in usual activities. Beyond this change between 2005 and 2008, the general pattern of the age trajectory remains almost unchanged over time. In particular, and especially for the severe limitation, we observed less rapid increase of the prevalence around the retirement age. Among men and women, the age standardized prevalence of reported disability increases over time (Table).

Standardized prevalence of activity limitation at age 15 and over (in %), SILC EU28, 2005, 2008, 2011 and 2014

The standardized rate of disability varies little over the years, even between 2005 and 2008 (period of changes in the instrument in some MS). Overall, these rates disclose a small increase over time in the prevalence of reported disability across the European Union.

	2005	2008	2011	2014
Men	23,0	23,1	23,0	23,9
(3-year gap)		(0,1)	(-0,1)	(0,8)
Women	27,8	28,2	28,6	29,3
(3-year gap)		(0,3)	(0,4)	(0,7)

BRIDGE-Health (Bridging Information and Data Generation for Evidence-based Health Policy and Research)

The **European Health and Life Expectancy Information System (EHLEIS)** is part of **BRIDGE-Health** which aims to prepare the transition towards a sustainable and integrated EU health information system within the third EU Health Program, 2014-2020 (www.bridge-health.eu).

